

SEE REVERSE SIDE OF BAG



**ORCHARD AGRI
CHEMICALS**

COPPER SULPHUR

Reg. No. L3564 Act/Wet No. 36 of/van 1947

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

GROUP

M

2

FUNGICIDES

A dusting powder for the control of plant diseases and mites as listed.

’n Stuifpoeier teen plantsiektes en myte soos aangedui.

WARNING



HAZARD STATEMENTS:

Flammable solid.
Causes skin irritation
Very toxic to aquatic life with long lasting effect

PRECAUTIONARY STATEMENTS:

Keep away from heat, sparks, open flames or hot surfaces - No Smoking. Ground and bond container and receiving equipment
Wear gloves, protective clothing, eye protection & face protection. Wash hands and face after handling.

ACTIVE INGREDIENTS / AKTIEWE BESTANDELE

Sulphur	980g/kg	Swael
Copper Oxychloride	850g/kg	Koperoksiechloried

In case of poisoning call the following number: 0861 555 777
Emergency number: 023 347 2754

25 kg

Batch Number

Lot Nommer :

Date of manufacture

Vervaardigingsdatum :

Registration holder/Registrasiehouer:

Orchard Agrikem (Pty) Ltd
Co. Reg. No./Mpy. Reg. Nr. 2014/196976/07

P.O. Box 94
Worcester, 6849

Tel: +27 (0)23 347 2754
www.orchardagri.co.za

UN No. 1350

COPPER SULPHUR

COPPER SULPHUR

WARNINGS

- **Flammable solid.**
- **Causes skin irritation**
- **Very toxic to fish with long lasting effects**
- **Keep out of reach of children**
- **may form combustible concentrations in air**
- **minimum number of days which must elaps between application and harvesting:**
Tomatoes, cucumbers - 3 days Table Grapes - 3 days Wine Grapes - 4 to 6 weeks
- **Re-entry: do not enter treated area within 1 day unless wearing protected clothing**

DISCLAIMER

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that this product will be efficacious under all conditions because the action and the effect thereof may be affected by factors such as abnormal climatic and storage conditions, compatibility with other substances not indicated on the label and occurrence of resistance of pests or disease to the remedy concerned, as well as by the method, time and accuracy of application.

The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
- Ground and bond container and receiving equipment.
- Wash hands and face thoroughly after handling.
- Avoid release to the environment.
- Wear protective gloves/protective colthing/eye protection/ face protection/hearing protection.
- IF ON SKIN: wash with plenty pf water.
- If skin irritation occurs: get medical help.
- Take off contaminated clothing and wash it before reuse.
- Collect spillage.

FIRST AID TREATMENT

Rinse immediately with plenty of water and treat symptomatically.

Relevant Substance

Component	Percentage
Sulphur	90 - 97%
Copper oxichloride	1 - 10%

APPLICATION RATES

Crop, Pest & Disease	Dosage Rate	Remarks
Tomatoes Early and late blight. Septoria leafspot.	15 to 30 kg / ha	In winter rainfall areas start dusting when the plants reach a height of 15 cm. In summer rainfall areas start dusting when disease threatens. Will also control rustmites and Erinos mite. Dosage rate depends on plant size.
Roses Powdery Mildew and Black Spot.	15 to 30 kg / ha	Apply at first sign of disease. Repeat weekly.
Other flowers and ornamentals Powdery Mildew and Downy Mildew.	10 to 30 kg / ha	Apply at first sign of disease. Will also control various leafspot and rust diseases.
Grapevines Anthracnose, Bacterial Blight, Dead Arm, Downy Mildew, Powdery Mildew(Oidium), Botrytis Rot and Erinos Mite.	15 to 30 kg / ha	Dust when shoots are 5 to 10 cm long and again when 25 cm long, just flowering and again 3 weeks later. More frequent treatment may be necessary in some districts.
Cucurbits (Except muskmelon and cucumbers.) Downy Mildew and Powdery Mildew	10 to 30 kg / ha	Dust at first sign of disease. Ensure coverage of both upper and lower leaf surface. Dosage depends on plant size.
Mangoes Powdery Mildew	20 to 40 kg / ha	Dust at first sign of infection and at full bloom. Repeat if blossoming is irregular and after heavy rain.