

SEE REVERSE SIDE OF BAG



DUSTING SULPHUR

Reg. No. L9127 Act/Wet No. 36 of/van 1947

READ LABEL BEFORE USE

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

GROUP	M	2	FUNGICIDES
-------	---	---	------------

A fungicide, acaricide and insecticide for agricultural use in the form of a dusting powder for the control of Powdery Mildew, Mites and Thrips on crops as mentioned.

‘n Swam-, myt- en insekdoder vir landboukundige gebruik in die vorm van ‘n stuifpoeier vir die beheer van Poeieragtige Skimmel, Myte en Blaaspootjies op gewasse soos aangedui.

WARNING



HAZARD STATEMENTS:
Flammable solid.
May cause skin irritation.

PRECAUTIONARY STATEMENTS:
Keep away from heat, sparks, open flames or hot surfaces - No Smoking.
Wear gloves, protective clothing, eye protection & face protection.

ACTIVE INGREDIENT / AKTIEWE BESTANDEEL

Sulphur 980 g/kg Swael

In case of poisoning call the following number: 0861 555 777
Emergency number: 023 347 2754

25 kg

Batch Number
Lot Nommer :

Date of manufacture
Vervaardigingsdatum :

Registration holder/Registrasiehouer:
Orchard Agri (Pty) Ltd
Co. Reg. No./Mpy. Reg. Nr. 2004/007007/07

P.O. Box 94
Worcester, 6849

Tel: +27 (0)23 347 2754
www.orchardagri.co.za

UN No. 1350

DUSTING SULPHUR

WARNINGS

- Flammable solid.
- Causes skin irritation.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, animals and uninformed persons.
- May form combustible dust concentrations in air.
- Wine grapes: allow 4 to 6 weeks between last application and harvest of wine grapes. Can cause leaf –burn when high temperatures prevail.
- Re-entry: Do not enter treated area within 1 day of application unless wearing protective clothing.

DISCLAIMER

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that this product will be efficacious under all conditions because the action and the effect thereof may be affected by factors such as abnormal climatic and storage conditions, compatibility with other substances not indicated on the label and occurrence of resistance of pests or disease to the remedy concerned, as well as by the method, time and accuracy of application.

The registration holder further does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or lack of performance of the remedy concerned, due to failure of the user to follow the label instructions or to the occurrence of conditions, which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Keep away from heat, hot surfaces, sparks, open flames and other ignition sources. No smoking.
- Ground and bond container and receiving equipment.
- Use explosion-proof electrical/ventilating/lighting equipment.
- Wash hands and face thoroughly after handling.
- Wear protective gloves/protective clothing/eye protection/face protection/hearing protection.
- IF ON SKIN: wash with plenty of water.
- If skin irritation occurs: get medical help.
- Take off contaminated clothing and wash it before reuse.
- In case of fire: use suitable material to extinguish.

FIRST AID TREATMENT

Rinse immediately with plenty of water and treat symptomatically.

DIRECTIONS FOR USE: Use only as directed.

Dosage depends on plant size and spacing.

RELEVANT SUBSTANCE

Component	Percentage
Sulphur (Active)	98%

APPLICATION RATES

Crop / Pest & Disease	Dosage Rate	Remarks
Avocados Thrips	10 to 40 kg / ha	Dust when pest is noticed.
Bananas Thrips & Flat mite	Light dusting	Can also be applied into the bags covering the bunches.
Beans Thrips	10 to 40 kg / ha	Full cover dusting. Repeat after 10 to 14 days.
Two-spotted spider mite (Red spider mite)	10 to 40 kg / ha	Dust as soon as the mites are noticed and repeat after 10 to 14 days, if necessary.
Cucurbits Powdery mildew	10 to 40 kg / ha	Do not use on cucumbers and muskmelons. Start dusting at first signs of this Mildew. Do not confuse with Downy mildew, which cannot be controlled with sulphur. Repeat at 7 - to 14 -day intervals as long as weather favours disease.
Citrus Thrips	Allow 4 weeks safety period between a dusting and a spray oil treatment. 50 to 100 kg / ha	Dust at 100 % petal drop and repeat after 10 days, if necessary.
Silver mite	50 to 100 kg / ha	Apply in spring if russetting occurred the previous season.
Rust mite & Flat mite	50 to 100 kg / ha	Apply when the first blemishes are noted on the fruit during December to May. It is advisable to treat the whole orchard.
Grey mite	50 to 100 kg / ha	Dust during winter or at petal fall.
Grapevines Powdery mildew	15 to 30 kg / ha	Heavy applications may cause burn, especially on hot days. Wine grapes: Do not apply after the berries have reached pea size (4 to 6 weeks prior to harvesting) as this may adversely influence fermentation. Winter rainfall region: Start dusting when shoots are 10 cm long and repeat every 3 weeks as long as necessary.
Erinose-bud mite	15 to 30 kg / ha	Summer rainfall region: Start dusting towards the middle of summer, particularly if rainfall is low or when first symptoms are noticed. Apply when shoots are approximately 10 cm long and repeat when the vines are in flower and again about the first week in December.
Lupins Powdery mildew	15 to 20 kg / ha	Dust when symptoms are noticed and repeat at 7- to 14-day intervals, if necessary. May cause leaf burn when high temperatures prevail.
Mangoes Powdery mildew	20 to 40 kg / ha	Dust at first sign of infection and at full blossom. Repeat if blossoming is irregular or if heavy rain falls.
Ornamentals Powdery mildew	10 to 40 kg / ha	Sulphur may injure certain species or cultivars. Dust every 10 to 14 days, depending on weather conditions favouring the disease. Ensure light but thorough coverage. In the case of roses dust the new growth weekly.
Thrips	10 to 40 kg / ha	Dust evenly at weekly intervals as soon as the insects or their damage is noticed.
Peaches Powdery mildew	20 to 40 kg / ha	Dust at first signs of infection and repeat every 10 to 14 days as long as the weather favours the disease.
Peas Powdery mildew	10 to 40 kg / ha	Apply to give an even light deposit, as soon as the first symptoms appear and repeat every 7 to 10 days as long as weather favours disease.
Tomatoes Erinose & Rust mite	10 to 40 kg / ha	Dust at first symptoms and repeat at 2- to 4-week intervals, if necessary.

DUSTING SULPHUR