

PRODUCT INFORMATION

Sluggit 30 Prima

Effective molluscicide

Sluggit Prima slug and snail bait with **30g/kg Metaldehyde** will ensure effective control for all slugs and snails.

Use Prima For Best Protection


All products in our Prima range is produced by Wet Extrusion, using advanced technology and superior ingredients to deliver a premium product.

- The baits are small and uniform in size
- Even spreading and higher density in bait distribution lead to better control of snails and slugs.
- Premium ingredients and manufacturing process produce a bait that is high in density, is robust and has better resistance to rain and elements, lasting up to **21 days** after application.
- Uniformity and robustness enhances mechanical spreading due to good ballistic properties.
- Unique formulation ensures a bait that is highly palatable to snails and slugs.
- Baits have the ability to soften quickly after contact with moisture, promoting fast and easy consumption by snails without compromising bait structure and longevity of product.
- The product is almost dust free.



Specifications

Appearance	Green pellets
Average Bait Size	2.2mm diam, 2.2 mm long
Baits/kg	More than 100 000
Metaldehyde Content	30g/kg
Available Pack Size	25kg 500kg 1 000kg
Process	Wet Extrusion
Shelf Life	2 years



PRODUCT INFORMATION



Withholding Periods

Harvest

Horticultural Crops/Vineyards

Do not apply later than 7 days before harvest.

Agricultural Crops

Do not apply later than 8 weeks before harvest.

Pastures/Pulses/Oil Seeds/Winter Cereals

Do not graze or cut for stock food for 5 weeks after application

Continue to monitor the crop and population numbers after applications and make follow up applications if necessary.

Snail monitoring to determine application times and rates

The key to efficient crop protection is continuous monitoring. Structured snail monitoring practices before, and 7 days after application should always be followed to determine snail population and appropriate application rates for effective control. Application rates should be seen as guidelines only and should be adjusted to address your specific snail population based on monitoring results.

Baits are seen as a source of food

Snails, slugs and insects see Sluggit baits as food. When they move around looking for food and happen to cross path with a bait, they will feed on it with toxic consequences. They will, at the same time, also feed on natural food sources. The more baiting points on the ground, the higher the likelihood of them feeding on it.

Protection period

Baits that last longer in the field, results in a longer protection period. Weather conditions have a big influence.

Why number of baits/kg is relevant

More baits/kg means more baiting points. This will lead to a higher likelihood of snails and slugs coming across baits in the field. Use 4 baits per slug or snail as a dosage guide. Sluggit 30 Prima has more than 100 000 baits per kg.

Timing is crucial

Apply during warmer, moist conditions just after rain when slugs and snails are more likely to feed. During Autumn, Spring and early Summer snails and slugs are known to be very active.

Application Rates

Situation	Application Rates
Agricultural Crops Cereals, oil seeds (including poppies) Pulses	6kg-15kg/Ha
Horticultural Crops Fruit, grapevines Vegetables	6kg-15kg/Ha
Pasture Establishment	5kg-8kg/Ha